arXiv:1212.5375v1 [math.CO] 21 Dec 2012

STRUCTURE COEFFICIENTS OF THE HECKE ALGEBRA OF $(\mathcal{S}_{2n}, \mathcal{B}_n)$

OMAR TOUT

Abstract. The Hecke algebra of the pair $(\mathcal{S}_{2n}, \mathcal{B}_n)$, where $\mathcal{B}_n$ is the hyperoctahedral subgroup of $\mathcal{S}_{2n}$, was introduced by James in 1961. It is a natural analogue of the center of the symmetric group algebra. In this paper, we give a polynomiality property of its structure coefficients. Our main tool is a combinatorial universal algebra which projects on the Hecke algebra of $(\mathcal{S}_{2n}, \mathcal{B}_n)$ for every n. To build it, we introduce new objects called partial bijections.

1. INTRODUCTION

The center of the symmetric group algebra of n, denoted $Z(\mathbb{C}[\mathcal{S}_n])$, is a classical object in algebraic combinatorics. It is linearly generated by elements $\mathcal{Z}_\lambda$, indexed by partitions of n, which are the sums of permutations of n with cycle-type λ. The structure coefficients $c_{\lambda\delta}^{\rho}$ describe the product in this algebra:

$$\mathcal{Z}_\lambda \mathcal{Z}_\delta = \sum_{\rho \text{ partition of } n} c_{\lambda\delta}^{\rho} \mathcal{Z}_\rho.$$

In other words, $c_{\lambda\delta}^{\rho}$ counts the number of pairs of permutations (x, y) with cycle-type λ and δ such that $x \cdot y = z$ for a fixed permutation z with cycle-type ρ. It is known, see [Cor75], that these coefficients also count numbers of graphs drawn on oriented surfaces with some additional conditions. One of the tools used to calculate these coefficients is the representation theory of the symmetric group, see [JV90, Lemma 3.3]. In [GS98, Theorem 2.1], Goupil and Schaeffer gave a cumbersome formula for $c_{\lambda\delta}^{\rho}$ if one of the partitions λ, δ and ρ is equal to (n). There are no formulas for $c_{\lambda\delta}^{\rho}$ in general.

In 1958, Farahat and Higman proved the polynomiality of the coefficients $c_{\lambda\delta}^{\rho}$ in n when λ, δ and ρ are fixed partitions, completed with parts equal to 1 to get partitions of n, see [FH59, Theorem 2.2]. This result is also proved by Ivanov and Kerov in [IK99] through the introduction of partial permutations. This proof provides a combinatorial description of these coefficients.

Here, we consider the Hecke algebra of the pair $(\mathcal{S}_{2n}, \mathcal{B}_n)$, where $\mathcal{B}_n$ is the hyperoctahedral group (see definition in section 2.2). It was introduced by James in [Jam61] and it also has a basis indexed by partitions of n. This algebra is a natural analogue of $Z(\mathbb{C}[\mathcal{S}_n])$ for several reasons. Goulden and Jackson proved in [GJ96] that its structure coefficients count graphs drawn on non-oriented surfaces. To get formulas for these coefficients, zonal characters are used instead of irreducible characters of the symmetric group, see [Mac95, Section VII, 2].

In this paper we give a polynomiality property of the structure coefficients of the Hecke algebra of $(\mathcal{S}_{2n}, \mathcal{B}_n)$. We prove that these coefficients can be written as the product of the number $2^n n!$ with a polynomial in n. In some specific basis, this polynomial has non-negative coefficients. Our proof is inspired by the construction of Ivanov and Kerov in [IK99]. However, we had to face some difficulties that do not appear in their work. In the proof, we introduce new combinatorial objects called partial bijections of n. These objects allow us to build in a combinatorial way a universal algebra which projects

---

1991 *Mathematics Subject Classification.* 05E15.

Key words and phrases. Hecke algebra of $(\mathcal{S}_{2n}, \mathcal{B}_n)$, partial bijections, structure coefficients.

Partially supported by ANR grant PSYCO ANR-11-JS02-001.

on Hecke algebra of $(\mathcal{S}_{2n}, \mathcal{B}_n)$ for every n. It also gives us a combinatorial description of the coefficients of the relevant polynomials.

A weaker version of our polynomiality result (without non-negativity of the coefficients) for the structure coefficients of Hecke algebra of $(\mathcal{S}_{2n}, \mathcal{B}_n)$ has been established by indirect approach using Jack polynomials in [DF12, Proposition 4.4]. There is no combinatorial description in that proof. In [AC10], Aker and Can considered the same question, but their article contains a mistake (the coefficient $2^n n!$ does not appear in their result).

The paper is organized as follows. In section 2, we put on all necessary definitions to describe the Hecke algebra of $(\mathcal{S}_{2n}, \mathcal{B}_n)$. Then, we give our main result about its structure coefficients. We start section 3 by introducing partial bijections of n then we construct our universal algebra. We use this algebra in section 4 to prove Theorem 2.1.

2. DEFINITIONS AND STATEMENT OF THE MAIN RESULT

2.1. **Partitions.** Since partitions index bases of the algebras studied in this paper, we recall the main definitions. A *partition* λ is a sequence of integers $(\lambda_1, \lambda_2, \dots)$ where $\lambda_1 \geq \lambda_2 \geq \dots \geq 1$. The λ_i are called the *parts* of λ and the *size* of λ, denoted by $|\lambda|$, is the sum of all its parts. If $|\lambda| = n$, we say that λ is a partition of n. We will also use the exponential notation $\lambda = (1^{m_1(\lambda)}, 2^{m_2(\lambda)}, 3^{m_3(\lambda)}, \dots)$, where $m_i(\lambda)$ is the number of parts equal to i in the partition λ. If λ and δ are two partitions we define the *union* $\lambda \cup \delta$ as the following partition:

$$\lambda \cup \delta = (1^{m_1(\lambda)+m_1(\delta)}, 2^{m_2(\lambda)+m_2(\delta)}, 3^{m_3(\lambda)+m_3(\delta)}, \dots).$$

A partition is called *proper* if it does not have any part equal to 1. The proper partition associated to a partition λ is the partition $\bar{\lambda} := \lambda \setminus (1^{m_1(\lambda)}) = (2^{m_2(\lambda)}, 3^{m_3(\lambda)}, \dots)$.

2.2. **Permutations and Coset type.** We will denote by $[n]$ the set $\{1, \dots, n\}$. A *permutation* of n is a bijection between the set $[n]$ and itself. For a permutation ω, we use the line notation $\omega_1\, \omega_2\, \cdots\, \omega_n$, where $\omega_i = \omega(i)$. The set $\mathcal{S}_n$ of all permutations of n is a group for the composition called the *symmetric group* of size n.

To each permutation ω of $2n$ we associate a graph $\Gamma(\omega)$ with $2n$ vertices located on a circle. Each vertex is labelled by two labels (exterior and interior). The exterior labels run through natural numbers from 1 to $2n$ around the circle. The interior label of the vertex with exterior label i is $\omega(i)$. We link the vertices with exterior (resp. interior) labels $2i-1$ and $2i$ by an exterior (resp. interior) edge. Since exterior and interior edges alternate, the graph $\Gamma(\omega)$ has only cycles of even lengths $2\lambda_1 \geq 2\lambda_2 \geq 2\lambda_3 \geq \cdots$. The *coset-type* of ω denoted by $ct(\omega)$ is the partition $(\lambda_1, \lambda_2, \lambda_3, \dots)$ of n.

Example 2.1. *The graph $\Gamma(\omega)$ associated to the permutation $\omega = 2\,4\,9\,3\,1\,10\,5\,8\,6\,7 \in \mathcal{S}_{2n}$ is drawn on Figure 1. It has two cycles of length 6 and 4, so $ct(\omega) = (3, 2)$.*

For every $k \geq 1$, we set $\rho(k) := \{2k-1, 2k\}$. The *hyperoctahedral group* $\mathcal{B}_n$ is the subgroup of $\mathcal{S}_{2n}$ of permutations ω such that, for every $1 \leq k \leq n,$ there exists $1 \leq k' \leq n$ with $\omega(\rho(k)) = \rho(k')$. In other words $\mathcal{B}_n = \{\omega \in \mathcal{S}_{2n} \mid ct(\omega) = (1^n)\}$. For example, $4\,3\,1\,2\,6\,5 \in \mathcal{B}_3$.

A $\mathcal{B}_n$*-double coset* of $\mathcal{S}_{2n}$ is the set $\mathcal{B}_n x \mathcal{B}_n = \{bxb' \ ; \ b, b' \in \mathcal{B}_n\}$ for some $x \in \mathcal{S}_{2n}$. It is known, see [Mac95, page 401], that two permutations of $\mathcal{S}_{2n}$ are in the same $\mathcal{B}_n$-double coset if and only if they have the same coset-type. Thus, if $x \in \mathcal{S}_{2n}$ has coset-type λ, we have:

$$\mathcal{B}_n x \mathcal{B}_n = \{y \in \mathcal{S}_{2n} \text{ such that } ct(y) = \lambda\}.$$



FIGURE 1. The graph $\Gamma(\omega)$ from Example 2.1.

2.3. **The Hecke algebra of $(\mathcal{S}_{2n}, \mathcal{B}_n)$.** The *symmetric group algebra of* n denoted by $\mathbb{C}[\mathcal{S}_n]$ is the algebra over $\mathbb{C}$ linearly generated by all permutations of n. The group $\mathcal{B}_n \times \mathcal{B}_n$ acts on $\mathbb{C}([\mathcal{S}_{2n}])$ by the following action: $(b, b') \cdot x = bxb'$, called the $\mathcal{B}_n$-double action. The *Hecke algebra* of $(\mathcal{S}_{2n}, \mathcal{B}_n)$ denoted by $\mathbb{C}[\mathcal{B}_n \setminus \mathcal{S}_{2n}/\mathcal{B}_n]$ is the sub-algebra of $\mathbb{C}[\mathcal{S}_{2n}]$ of elements invariant under the $\mathcal{B}_n$-double action. Recall that $\mathcal{B}_n$-double cosets are indexed by partitions of n. Therefore, the set

$$\{K_\lambda(n) : \lambda \text{ proper partition, } |\lambda| \leq n\}$$

forms a basis for $\mathbb{C}[\mathcal{B}_n \setminus \mathcal{S}_{2n}/\mathcal{B}_n]$, where $K_\lambda(n)$ is the sum of all permutations of $\mathcal{S}_{2n}$ with coset-type $\lambda \cup 1^{n-|\lambda|}$. So, for any two proper partitions λ and δ with size at most n, there exist complex numbers $\alpha^\rho_{\lambda\delta}(n)$ such that:

$$K_\lambda(n) \cdot K_\delta(n) = \sum_{\substack{\rho \text{ proper partition} \\ |\rho| \leq n}} \alpha^\rho_{\lambda\delta}(n) K_\rho(n). \tag{1}$$

2.4. **Main result.** We give in this paper a polynomiality property of the structure coefficients of the Hecke algebra of $(\mathcal{S}_{2n}, \mathcal{B}_n)$ that appear in (1). More precisely, we prove the following theorem.

Theorem 2.1. *Let λ, δ and ρ be three proper partitions, we have:*

$$\alpha^\rho_{\lambda\delta}(n) = \begin{cases} 2^n n! f^\rho_{\lambda\delta}(n) & \text{if} \quad n \geq |\rho|, \\ 0 & \text{if} \quad n < |\rho|, \end{cases}$$

where $f^\rho_{\lambda\delta}(n) = \sum_{j=0}^{|\lambda|+|\delta|-|\rho|} a_j (n - |\rho|)_j$ is a polynomial in n with $a_j \in \mathbb{Q}^+$.

Example 2.2. *Let us compute the structure coefficient $\alpha^{\emptyset}_{(2)(2)}(n)$. We have:*

$$K_{(2)}(n) = \sum_{\substack{\omega \in \mathcal{S}_{2n} \\ ct(\omega) = (2) \cup (1^{n-2})}} \omega.$$

To find the coefficient of $K_\emptyset(n)$ in $K_{(2)}(n) \cdot K_{(2)}(n)$, we fix a permutation with coset-type (1^n), for example Id_{2n}, and we look in how many ways we can obtain Id_{2n} as a product of two elements $\sigma \cdot \beta$ where $ct(\sigma) = ct(\beta) = (2, 1^{n-2})$. Thus we are looking for the number of permutations $\sigma \in \mathcal{S}_{2n}$ such that $ct(\sigma) = ct(\sigma^{-1}) = (2, 1^{n-2})$. But, for any $\sigma \in \mathcal{S}_{2n}$ with $ct(\sigma) = (2, 1^{n-2})$, his inverse has the *same coset-type. Therefore* $\alpha^{\emptyset}_{(2)(2)}(n)$ *is the number of permutations of coset-type* $(2,1^{n-2})$, *which is by* [Mac95, page 402]

$$\frac{(2^n n!)^2}{2^{n-1}(2\cdot(n-2)!)}=n(n-1)2^n n!.$$

2.5. **Major steps of the proof.** The idea of the proof is to build a universal algebra $\mathcal{A}_\infty$ over $\mathbb{C}$ satisfying the following properties:

(1) For every $n\in\mathbb{N}^*$, there exists a morphism of algebras $\theta_n:\mathcal{A}_\infty\longrightarrow\mathbb{C}[\mathcal{B}_n\setminus\mathcal{S}_{2n}/\mathcal{B}_n]$.

(2) Every element x in $\mathcal{A}_\infty$ is written in a unique way as an infinite linear combination of elements T_λ, indexed by partitions. For any two partitions λ and δ, there exist non-negative rational numbers $b^{\rho}_{\lambda\delta}$ such that:

$$T_\lambda * T_\delta=\sum_{\rho \text{ partition}} b^{\rho}_{\lambda\delta}T_\rho. \tag{2}$$

(3) The morphism θ_n sends T_λ to a multiple of $K_{\bar{\lambda}}(n)$.

To build $\mathcal{A}_\infty$, we introduce new combinatorial objects called *partial bijections*. For every $n\in\mathbb{N}^*$, we construct an algebra $\mathcal{A}_n$ using the set of partial bijections of size n. The algebra $\mathcal{A}_\infty$ is defined as the projective limit of this sequence $(\mathcal{A}_n)$.

The projection $p_n:\mathcal{A}_\infty\to\mathcal{A}_n$ involves coefficients which are polynomials in n. By defining the extension of a partial bijection of n to the set $[2n]$, we construct a morphism from $\mathcal{A}_n$ to $\mathbb{C}[\mathcal{B}_n\setminus\mathcal{S}_{2n}/\mathcal{B}_n]$. Its coefficients involve the number $2^n n!$. It turns out that the morphism θ_n is the composition of those two morphisms:

$$\begin{tikzcd}
\mathcal{A}_\infty \arrow[dr, "p_n"] \arrow[dd, "\theta_n"'] & \\
& \mathcal{A}_n \arrow[dl] \\
\mathbb{C}[\mathcal{B}_n\setminus\mathcal{S}_{2n}/\mathcal{B}_n] &
\end{tikzcd}$$

The final step consists in applying the chain of morphisms in the diagram above to equation (2).

Remark. This method is based on Ivanov and Kerov's one to get the polynomiality of the structure coefficients of the center of the symmetric group algebra (see [IK99] for more details). Nevertheless, our construction is more complicated, mainly because a partial bijection does not have a unique trivial extension to a given set, see Definition 3.2.

3. THE PARTIAL BIJECTION ALGEBRA

In this section we define the set of partial bijections of n. With this set, we build the algebras and morphisms that appear in the diagram above.

3.1. **Definition.** We start by defining partial bijections of n and the partial bijection algebra. Then, we introduce the notion of trivial extension of a partial bijection of n and we use it to build a morphism between the partial bijection algebra of n and the symmetric group algebra of $2n$.

For $n\in\mathbb{N}^*$, $\mathbf{P}_n$ denotes the following set:

$$\mathbf{P}_n:=\{\rho(k_1)\cup\cdots\cup\rho(k_i)\mid 1\le i\le n,\ 1\le k_1<\cdots<k_i\le n\}.$$

For each $1\le k\le n$, the number $\frac{n!}{(n-k)!}=n(n-1)\cdots(n-k+1)$ is denoted by $(n)_k$.

Definition 3.1. A *partial bijection* of n is a triple $(\sigma, d, d^{'})$ where $d, d^{'} \in \mathbf{P}_n$ and $\sigma : d \longrightarrow d^{'}$ is a bijection. We denote by Q_n the set of all partial bijections of n.

It should be clear that

$$|Q_n| = s_n = \sum_{k=0}^{n} \binom{n}{k}^2 (2k)!.$$

A permutation σ of $2n$ can be written as $(\sigma, [2n], [2n])$, so the set $\mathcal{S}_{2n}$ can be considered as a subset of Q_n.
For any partial bijection α, we will use the convention that σ (resp. d, $d^{'}$) is the first (resp. second, third) element of the triple defining α. The same convention holds for $\widetilde{\alpha}$, α_i, $\hat{\alpha} \dots$

Observation. In the same way as in section 2.2, we can associate to each partial bijection α of n a graph $\Gamma(\alpha)$ with $|d|$ vertices placed on a circle. The exterior (resp. interior) labels are the elements of the set d (resp. $d^{'}$). Since the sets d and $d^{'}$ are in $\mathbf{P}_n$, we can link $2i$ with $2i-1$ as in the case $d = d^{'} = [2n]$. So, the definition of coset-type extends naturally to partial bijection. We denote by $ct(\alpha)$ or $ct(\sigma)$ the coset-type of a partial bijection α.

Definition 3.2. Let $(\sigma, d, d^{'})$ and $(\widetilde{\sigma}, \widetilde{d}, \widetilde{d^{'}})$ be two partial bijections of n. We say that $(\widetilde{\sigma}, \widetilde{d}, \widetilde{d^{'}})$ is a *trivial extension* of $(\sigma, d, d^{'})$ if:

$$d \subseteq \widetilde{d},\ \widetilde{\sigma}_{|_d} = \sigma \text{ and } ct(\widetilde{\sigma}) = ct(\sigma) \cup \left(1^{\frac{|\widetilde{d} \setminus d|}{2}}\right).$$

We denote by $P_\alpha(n)$ the set of all trivial extensions of α in Q_n.

Lemma 3.1. *Let α be a partial bijection of n and X an element of $\mathbf{P}_n$ such that $d \subseteq X$. The number of trivial extensions $\widetilde{\alpha}$ such that $\widetilde{d} = X$ is*

$$(2n - |d|) \cdot (2n - |d| - 2) \cdots (2n - |d| - |X \setminus d| + 2) = 2^{\frac{|X \setminus d|}{2}} \left(n - \frac{|d|}{2}\right)_{\frac{|X \setminus d|}{2}}.$$

We have the same formula for the number of trivial extensions $\widetilde{\alpha}$ such that $\widetilde{d'} = X$.

Proof. We proceed by induction on the size of $X \setminus d$. If $|X \setminus d| = 2$, suppose that $X \setminus d = \rho(k)$ for some $k \in [n]$. There are $2n - |d|$ possible values for $\widetilde{\sigma}(2k-1)$. If $\widetilde{\sigma}(2k-1) = 2k'-1$ (resp. $2k'$), we have $\widetilde{\sigma}(2k) = 2k'$ (resp. $2k'-1$). So, the number of trivial extensions $\widetilde{\alpha}$ such that $\widetilde{d} = X$ is $2n - |d|$.
We suppose that we have the result for $|X \setminus d| \leq 2(r-1)$. Let X be a set such that $|X \setminus d| = 2r$. We fix an element $2i-1$ of $X \setminus d$. Trivial extensions $\widetilde{\alpha}$ such that $\widetilde{d} = X$ are obtained as follows:

- first, take all trivial extensions α_1 such that $d_1 = X \setminus \rho(i)$. Since $\big|\big(X \setminus \rho(i)\big) \setminus d\big| = 2(r-1)$, the number of these trivial extensions is by induction $2^{r-1}(n - \frac{|d|}{2})_{r-1}$.
- second, for every trivial extension α_1, take all trivial extensions $\widetilde{\alpha_1}$ such that $\widetilde{d_1} = X$. For a fixed α_1, the number of trivial extensions $\widetilde{\alpha_1}$ such that $\widetilde{d_1} = X$ is, by the base case of induction, $2n - |X \setminus \rho(i)| = 2n - |d| - 2r + 2$.

Every trivial extension $\widetilde{\alpha}$ is obtained exactly once: as a trivial extension of α_1, where α_1 is $\widetilde{\alpha}_{|_{X \setminus \rho(i)}}$. Thus, the number of trivial extension $\widetilde{\alpha}$ such that $\widetilde{d} = X$ is the product:

$$2^{r-1}\left(n - \frac{|d|}{2}\right)_{r-1} \cdot (2n - |d| - 2r + 2) = 2^r \left(n - \frac{|d|}{2}\right)_r.$$

This ends our induction and proves the first part of lemma. The proof of the second part (number of trivial extension $\widetilde{\alpha}$ such that $\widetilde{d'} = X$) is similar. $\square$

Consider $\mathcal{D}_n = \mathbb{C}[Q_n]$ the vector space with basis Q_n. We want to endow it with an algebra structure. Let α_1 and α_2 be two partial bijections. If $d_1 = d'_2$, we can compose α_1 and α_2 and we define $\alpha_1 * \alpha_2 = \alpha_1 \circ \alpha_2 = (\sigma_1 \circ \sigma_2, d_2, d'_1)$. Otherwise, we need to extend α_1 and α_2 to partial bijections $\widetilde{\alpha_1}$ and $\widetilde{\alpha_2}$ such that $\widetilde{d_1} = \widetilde{d'_2}$. Since there exist several trivial extensions of α_1 and α_2, a natural choice is to take the average of the composition of all possible trivial extensions. Let $E^{\alpha_2}_{\alpha_1}(n)$ be the following set:

$$E^{\alpha_2}_{\alpha_1}(n) \quad := \quad \{(\widetilde{\alpha_1}, \widetilde{\alpha_2}) \in P_{\alpha_1}(n) \times P_{\alpha_2}(n) \text{ such that } \widetilde{d_1} = \widetilde{d'_2} = d_1 \cup d'_2\}.$$

Elements of $E^{\alpha_2}_{\alpha_1}(n)$ are schematically represented on Figure 2.

Throughout the paper, we will use the following conventions on figures that represent elements of some sets.

- The data defining the set (so fixed when we go from an element of the set to another) is drawn using plain shapes.
- The element of the set is drawn using dashed shapes.

We define the product of α_1 and α_2 as follows:

$$\alpha_1 * \alpha_2 := \frac{1}{|E^{\alpha_2}_{\alpha_1}(n)|} \sum_{(\widetilde{\alpha_1}, \widetilde{\alpha_2}) \in E^{\alpha_2}_{\alpha_1}(n)} \widetilde{\alpha_1} \circ \widetilde{\alpha_2}. \tag{3}$$

By Lemma 3.1, we can see that:

$$|E^{\alpha_2}_{\alpha_1}(n)| = 2^{\frac{|d'_2 \setminus d_1|}{2} + \frac{|d_1 \setminus d'_2|}{2}} . (n - \frac{|d'_1|}{2})_{(\frac{|d'_2 \setminus d_1|}{2})} . (n - \frac{|d_2|}{2})_{(\frac{|d_1 \setminus d'_2|}{2})}. \tag{4}$$



FIGURE 2. Schematic representation of elements of $E^{\alpha_2}_{\alpha_1}(n)$.

Proposition 3.2. *The product $*$ is associative. In other words, $\mathcal{D}_n$ is a (non-unital) algebra.*

Proof. Postponed to next section. ☐

Proposition 3.3. *The following function defines a morphism of algebras:*

$$\begin{array}{rccc} \psi_n & : & \mathbb{C}[Q_n] & \to & \mathbb{C}[\mathcal{S}_{2n}] \\ & & \alpha & \mapsto & \frac{1}{2^{n-\frac{|d|}{2}}(n-\frac{|d|}{2})!} \displaystyle\sum_{\hat{\alpha} \in \mathcal{S}_{2n} \cap P_\alpha(n)} \hat{\sigma} \end{array} .$$

Proof. Let α_1 and α_2 be two basis elements of $\mathbb{C}[Q_n]$. We refer to Figure 2 and denote $2b = |d_2| = |d'_2|, 2c = |d_1| = |d'_1|$ and $2e = |d'_2 \cap d_1|$. We first prove that:

$$\begin{aligned} &\sum_{\widehat{\alpha_1} \in \mathcal{S}_{2n} \cap P_{\alpha_1}(n)} \sum_{\widehat{\alpha_2} \in \mathcal{S}_{2n} \cap P_{\alpha_2}(n)} \widehat{\sigma_1} \circ \widehat{\sigma_2} \\ &\qquad = 2^{n-(b+c-e)}(n-(b+c-e))! \sum_{(\widetilde{\alpha_1}, \widetilde{\alpha_2}) \in E^{\alpha_1}_{\alpha_2}(n)} \sum_{\widehat{\widetilde{\alpha_1} \circ \widetilde{\alpha_2}} \in \mathcal{S}_{2n} \cap P_{\widetilde{\alpha_1} \circ \widetilde{\alpha_2}}(n)} \widehat{\widetilde{\sigma_1} \circ \widetilde{\sigma_2}}. \end{aligned} \tag{5}$$

We fix $(\widetilde{\alpha_1}, \widetilde{\alpha_2}) \in E^{\alpha_2}_{\alpha_1}(n)$ and $\omega \in \mathcal{S}_{2n} \cap P_{\widetilde{\alpha_1}\circ\widetilde{\alpha_2}}(n)$, *i.e.*:

$$\omega_{|_{\widetilde{d_2}}} = \widetilde{\sigma_1} \circ \widetilde{\sigma_2} \text{ and } ct(\omega) = ct(\widetilde{\sigma_1} \circ \widetilde{\sigma_2}) \cup (1^{(n-(b+c-e))}).$$

We look for the number of permutations $\widehat{\sigma_1}$ and $\widehat{\sigma_2}$ in $\mathcal{S}_{2n} \cap P_{\alpha_1}(n)$ and $\mathcal{S}_{2n} \cap P_{\alpha_2}(n)$ such that $\widehat{\sigma_1} \circ \widehat{\sigma_2} = \omega$. In this equation, $\widehat{\sigma_2}$ determines $\widehat{\sigma_1}$. But the condition $\omega_{|_{\widetilde{d_2}}} = \widetilde{\sigma_1} \circ \widetilde{\sigma_2}$ gives the values of $\widehat{\sigma_2}$ on $\widetilde{d_2}$ ($\widehat{\sigma_2}(x) = \sigma_2(x)$ if $x \in d_2$ and $\widehat{\sigma_2}(x) = \sigma_1^{-1}(\omega(x))$ if $x \in \widetilde{d_2} \backslash d_2$). Thus, the number of ways to choose $\widehat{\sigma_2}$ is the number of ways to extend trivially $\widetilde{\sigma_2}$ to be a permutation of $2n$, which is $2^{n-(b+c-e)}(n-(b+c-e))!$ by Lemma 3.1. This proves equation (5).

Now we have:

$$\psi_n(\alpha_1)\psi_n(\alpha_2) = \frac{1}{2^{2n-b-c}(n-c)!(n-b)!} \sum_{\widehat{\alpha_1} \in \mathcal{S}_{2n} \cap P_{\alpha_1}(n)} \sum_{\widehat{\alpha_2} \in \mathcal{S}_{2n} \cap P_{\alpha_2}(n)} \widehat{\sigma_1} \circ \widehat{\sigma_2} \tag{6}$$

$$= \frac{(n-b-c+e)!}{2^{n-e}(n-c)!(n-b)!} \sum_{(\widetilde{\alpha_1}, \widetilde{\alpha_2}) \in E^{\alpha_1}_{\alpha_2}(n)} \sum_{\widehat{\widetilde{\alpha_1} \circ \widetilde{\alpha_2}} \in \mathcal{S}_{2n} \cap P_{\widetilde{\alpha_1} \circ \widetilde{\alpha_2}}(n)} \widehat{\widetilde{\sigma_1} \circ \widetilde{\sigma_2}}.$$

On the other hand:

$$\psi_n(\alpha_1 * \alpha_2) = \frac{1}{2^{b+c-2e}(n-c)_{(b-e)}(n-b)_{(c-e)}} \sum_{(\widetilde{\alpha_1}, \widetilde{\alpha_2}) \in E^{\alpha_1}_{\alpha_2}(n)} \psi_n\big((\widetilde{\sigma_1} \circ \widetilde{\sigma_2}, \widetilde{d_2}, \widetilde{d'_1})\big).$$

But

$$\psi_n\big((\widetilde{\sigma_1} \circ \widetilde{\sigma_2}, \widetilde{d_2}, \widetilde{d'_1})\big) = \frac{1}{2^{n-(b+c-e)}(n-(b+c-e))!} \sum_{\widehat{\widetilde{\alpha_1} \circ \widetilde{\alpha_2}} \in \mathcal{S}_{2n} \cap P_{\widetilde{\alpha_1} \circ \widetilde{\alpha_2}}(n)} \widehat{\widetilde{\sigma_1} \circ \widetilde{\sigma_2}}.$$

Thus,

$$\psi_n(\alpha_1 * \alpha_2) = \frac{(n-b-c+e))!}{2^{n-e}(n-c)!(n-b)!} \sum_{(\widetilde{\alpha_1}, \widetilde{\alpha_2}) \in E^{\alpha_1}_{\alpha_2}(n)} \sum_{\widehat{\widetilde{\alpha_1} \circ \widetilde{\alpha_2}} \in \mathcal{S}_{2n} \cap P_{\widetilde{\alpha_1} \circ \widetilde{\alpha_2}}(n)} \widehat{\widetilde{\sigma_1} \circ \widetilde{\sigma_2}}. \tag{7}$$

Comparing equations (6) and (7), we see that for any two partial bijections α_1 and α_2 of n, we have $\psi_n(\alpha_1 * \alpha_2) = \psi_n(\alpha_1)\psi_n(\alpha_2)$. In other words, ψ_n is a morphism of algebras. □

3.2. **Proof of the associativity of $*$.** Let α_1, α_2 and α_3 be three elements of Q_n. By definition of the product we have:

$$(\alpha_1 * \alpha_2) * \alpha_3 = \frac{1}{|E^{\alpha_2}_{\alpha_1}(n)|} \sum_{(\widetilde{\alpha_1}, \widetilde{\alpha_2}) \in E^{\alpha_2}_{\alpha_1}(n)} \frac{1}{|E^{\alpha_3}_{\widetilde{\alpha_1} \circ \widetilde{\alpha_2}}(n)|} \sum_{(\widetilde{\widetilde{\alpha_1} \circ \widetilde{\alpha_2}}, \widetilde{\alpha_3}) \in E^{\alpha_3}_{\widetilde{\alpha_1} \circ \widetilde{\alpha_2}}(n)} (\widetilde{\widetilde{\sigma_1} \circ \widetilde{\sigma_2}} \circ \widetilde{\sigma_3}, \widetilde{d_3}, \widetilde{\widetilde{d'_1}}), \tag{8}$$

$$\alpha_1 * (\alpha_2 * \alpha_3) = \frac{1}{|E^{\alpha_3}_{\alpha_2}(n)|} \sum_{(\widetilde{\alpha_2}, \widetilde{\alpha_3}) \in E^{\alpha_3}_{\alpha_2}(n)} \frac{1}{|E^{\widetilde{\alpha_2} \circ \widetilde{\alpha_3}}_{\alpha_1}(n)|} \sum_{(\widetilde{\alpha_1}, \widetilde{\widetilde{\alpha_2} \circ \widetilde{\alpha_3}}) \in E^{\widetilde{\alpha_2} \circ \widetilde{\alpha_3}}_{\alpha_1}(n)} (\widetilde{\sigma_1} \circ \widetilde{\widetilde{\sigma_2} \circ \widetilde{\sigma_3}}, \widetilde{\widetilde{d_3}}, \widetilde{d'_1}). \tag{9}$$

We consider now the following sets, indexing the double sums in the equations above:

$$X_1 = \{\big((\widetilde{\alpha_1}, \widetilde{\alpha_2}), (\widetilde{\widetilde{\alpha_1} \circ \widetilde{\alpha_2}}, \widetilde{\alpha_3})\big) \in E^{\alpha_2}_{\alpha_1}(n) \times E^{\alpha_3}_{\widetilde{\alpha_1} \circ \widetilde{\alpha_2}}(n)\},$$

and

$$X_2 = \{\big((\widetilde{\alpha_2}, \widetilde{\alpha_3}), (\widetilde{\alpha_1}, \widetilde{\widetilde{\alpha_2} \circ \widetilde{\alpha_3}})\big) \in E^{\alpha_3}_{\alpha_2}(n) \times E^{\widetilde{\alpha_2} \circ \widetilde{\alpha_3}}_{\alpha_1}(n)\}.$$

Using X_1, the product $(\alpha_1 * \alpha_2) * \alpha_3$ can be written as follows:

$$(\alpha_1 * \alpha_2) * \alpha_3 = \sum_{\big((\widetilde{\alpha_1},\widetilde{\alpha_2}),(\widetilde{\widetilde{\alpha_1}\circ\widetilde{\alpha_2}},\widetilde{\alpha_3})\big)\in X_1} \frac{1}{|E^{\alpha_2}_{\alpha_1}(n)|\cdot|E^{\alpha_3}_{\widetilde{\alpha_1}\circ\widetilde{\alpha_2}}(n)|}(\widetilde{\widetilde{\sigma_1}\circ\widetilde{\sigma_2}}\circ\widetilde{\sigma_3}, \widetilde{d_3}, \widetilde{\widetilde{d'_1}}). \tag{10}$$

Schematically, the elements in X_1 are represented on Figure 3.



FIGURE 3. Schematic representation of elements in X_1.

In the same way, using X_2, the product $\alpha_1 * (\alpha_2 * \alpha_3)$ can be written in the following way:

$$\alpha_1 * (\alpha_2 * \alpha_3) = \sum_{\big((\widetilde{\alpha_2},\widetilde{\alpha_3}),(\widetilde{\alpha_1},\widetilde{\widetilde{\alpha_2}\circ\widetilde{\alpha_3}})\big)\in X_2} \frac{1}{|E^{\alpha_3}_{\alpha_2}(n)|\cdot|E^{\widetilde{\alpha_2}\circ\widetilde{\alpha_3}}_{\alpha_1}(n)|}(\widetilde{\sigma_1}\circ\widetilde{\widetilde{\sigma_2}\circ\widetilde{\sigma_3}}, \widetilde{\widetilde{d_3}}, \widetilde{d'_1}). \tag{11}$$

Schematically, the elements in X_2 are represented on Figure 4. To prove the associativity of the product,



FIGURE 4. Schematic representation of elements of X_2.

we construct a set X and two surjective functions $\phi_1 : X \longrightarrow X_1$ and $\phi_2 : X \longrightarrow X_2$ in order to write both sums (equations (10) and (11)) as sums over the same set X. Let X be the set of elements

$$\big(\epsilon_1 = (\tau_1, \delta_0, \delta_1), \epsilon_2 = (\tau_2, \delta_1, \delta_2), \epsilon_3 = (\tau_3, \delta_2, \delta_3)\big) \in P_{\alpha_1}(n) \times P_{\alpha_2}(n) \times P_{\alpha_3}(n),$$

satisfying the following properties:

(1) $d_3^{'} \cup d_2 \subseteq \delta_2$.
(2) $d_2^{'} \cup d_1 \subseteq \delta_1$.
(3) $\delta_2 = d'_3 \cup \tau_2^{-1}(d_1 \cup d'_2)$.

The elements of this set are schematically represented on Figure 5. Note that (3) is a minimality condition. We will see in the proof of Lemma 3.4 below why it is useful.



FIGURE 5. Schematic representation of elements of X.

We define the maps $\phi_1 : X \longrightarrow X_1$ and $\phi_2 : X \longrightarrow X_2$ as follows:

$$\phi_1(\epsilon_1, \epsilon_2, \epsilon_3) = \big((\epsilon_{1_{|_{d_1 \cup d'_2}}}, \epsilon_{2_{|_{\tau_2^{-1}(d_1 \cup d'_2)}}}), (\epsilon_1 \circ \epsilon_2, \epsilon_3)\big),$$

$$\phi_2(\epsilon_1, \epsilon_2, \epsilon_3) = \big((\epsilon_{2_{|_{d_2 \cup d'_3}}}, \epsilon_{3_{|_{\tau_3^{-1}(d_2 \cup d'_3)}}}), (\epsilon_2 \circ \epsilon_3, \epsilon_1)\big).$$

The map ϕ_1 (resp. ϕ_2) forgets the dashed ellipse shape at the top (resp. bottom) of the third (resp. second) column of Figure 5. We denote by $2a = |d_3| = |d'_3|, 2b = |d_2| = |d'_2|, 2c = |d_1| = |d'_1|, 2d = |d_3^{'} \cap d_2|$ and $2e = |d_2^{'} \cap d_1|$. We prove the following lemma:

Lemma 3.4. *Informally, the map ϕ_1 is well defined and surjective. For any element $\big((\widetilde{\alpha_1}, \widetilde{\alpha_2}), (\widetilde{\widetilde{\alpha_1} \circ \widetilde{\alpha_2}}, \widetilde{\alpha_3})\big) \in X_1$, we have:*

$$|\phi_1^{-1}\Big(\big((\widetilde{\alpha_1}, \widetilde{\alpha_2}), (\widetilde{\widetilde{\alpha_1} \circ \widetilde{\alpha_2}}, \widetilde{\alpha_3})\big)\Big)| = 2^{a-d-f} \cdot (n-b-c+e)_{(a-d-f)},$$

where, $2f = |(\widetilde{d_2} \setminus d_2) \cap d_3^{'}|$.

Proof. For any $(\epsilon_1, \epsilon_2, \epsilon_3) \in X$, we must check that $\phi_1\big((\epsilon_1, \epsilon_2, \epsilon_3)\big)$ is in X_1. Set

$$(\widetilde{\alpha_1}, \widetilde{\alpha_2}) = (\epsilon_{1_{|_{d_1 \cup d'_2}}}, \epsilon_{2_{|_{\tau_2^{-1}(d_1 \cup d'_2)}}}).$$

First $(\widetilde{\alpha_1}, \widetilde{\alpha_2})$ is in $P_{\alpha_1}(n) \times P_{\alpha_2}(n)$ since (ϵ_1, ϵ_2) is in $P_{\alpha_1}(n) \times P_{\alpha_2}(n)$. Then, the arrival domain of $\widetilde{\alpha_2}$ and the definition domain of $\widetilde{\alpha_1}$ are $d_1 \cup d'_2$. Thus, $(\widetilde{\alpha_1}, \widetilde{\alpha_2})$ is in $E_{\alpha_1}^{\alpha_2}(n)$.

Second, it is easy to check that $(\epsilon_1 \circ \epsilon_2, \epsilon_3)$ is in $P_{\widetilde{\alpha_1} \circ \widetilde{\alpha_2}}(n) \times P_{\alpha_3}(n)$. Then, to be in $E^{\alpha_3}_{\widetilde{\alpha_1} \circ \widetilde{\alpha_2}}(n)$, $(\epsilon_1 \circ \epsilon_2, \epsilon_3)$ must verify the condition that $\delta_2 = d'_3 \cup \tau_2^{-1}(d_1 \cup d'_2)$, which is given by condition (3).

Thus, ϕ_1 is well defined.

Now, fix $\big((\widetilde{\alpha_1},\widetilde{\alpha_2}),(\widetilde{\widetilde{\alpha_1}\circ\widetilde{\alpha_2}},\widetilde{\alpha_3})\big) \in X_1$. We will count the number of its pre-images by ϕ_1. To construct an element of $\phi_1^{-1}\Big(\big((\widetilde{\alpha_1},\widetilde{\alpha_2}),(\widetilde{\widetilde{\alpha_1}\circ\widetilde{\alpha_2}},\widetilde{\alpha_3})\big)\Big)$, we have only to build ϵ_1,ϵ_2 and δ_1 since the other elements $\epsilon_3,\delta_0,\delta_2$ and δ_3 are determined by $\widetilde{\alpha_1},\widetilde{\alpha_2},\widetilde{\alpha_3}$ and $\widetilde{\widetilde{\alpha_1}\circ\widetilde{\alpha_2}}$. First, to build δ_1, we must extend $d_1\cup d_2^{'}$ by adding pairs of form $\rho(k)$ to obtain a set which has the same cardinality as $\delta_2=\widetilde{d_2}\cup d_3^{'}$ which is $2a+2b+2c-2e-2d-2f$ by condition. We have $|d_1\cup d_2^{'}|=2b+2c-2e$, so the number of possible ways to extend $d_1\cup d_2^{'}$ is the number of choice of $2a+2b+2c-2e-2d-2f-(2b+2c-2e)=2a-2d-2f$ elements among $2n-(2b+2c-2e)$. Since our choice must respect the condition that the extended set is in $\mathbf{P}_n$, this number is:

$$\binom{n-(b+c-e)}{a-d-f}.$$

Once the set $d_1\cup d_2^{'}$ is extended, we should extend $\widetilde{\sigma_1}$ to τ_1 (we have the definition domain δ_0 and the arrival domain δ_1) by sending the pairs of form $\rho(k)$ to pairs with same form. The number of ways to do this is:

$$2^{a-d-f}\cdot(a-d-f)!.$$

After extending $\widetilde{\sigma_1}$ to τ_1, we get immediately τ_2 because $\tau_1\circ\tau_2=\widetilde{\widetilde{\sigma_1}\circ\widetilde{\sigma_2}}$ is given. Thus, the cardinality of the set $\phi_1^{-1}\Big(\big((\widetilde{\alpha_1},\widetilde{\alpha_2}),(\widetilde{\widetilde{\alpha_1}\circ\widetilde{\alpha_2}},\widetilde{\alpha_3})\big)\Big)$ is equal to:

$$\binom{n-(b+c-e)}{a-d-f}\cdot 2^{a-d-f}\cdot(a-d-f)!=2^{a-d-f}\cdot(n-b-c+e)_{(a-d-f)}.$$

□

In the same way, we can prove the following lemma.

Lemma 3.5. *The map ϕ_2 is well defined and surjective. For any element $\big((\widetilde{\alpha_2},\widetilde{\alpha_3}),(\widetilde{\alpha_1},\widetilde{\widetilde{\alpha_2}\circ\widetilde{\alpha_3}})\big)\in X_2$, we have:*

$$|\phi_2^{-1}\Big(\big((\widetilde{\alpha_2},\widetilde{\alpha_3}),(\widetilde{\alpha_1},\widetilde{\widetilde{\alpha_2}\circ\widetilde{\alpha_3}})\big)\Big)|=2^{c-e-g}\cdot(n-a-b+d)_{(c-e-g)},$$

where, $2g=|(\widetilde{d_2^{'}}\setminus d_2^{'})\cap d_1|$.

We can also verify using the notations above that:

$$|E_{\alpha_1}^{\alpha_2}(n)|=2^{b+c-2e}\cdot(n-c)_{(b-e)}\cdot(n-b)_{(c-e)},$$

$$|E_{\alpha_2}^{\alpha_3}(n)|=2^{a+b-2d}\cdot(n-b)_{(a-d)}\cdot(n-a)_{(b-d)},$$

and

$$|E_{\widetilde{\alpha_1}\circ\widetilde{\alpha_2}}^{\alpha_3}(n)|=2^{a+b+c-2d-e-2f}\cdot(n-b-c+e)_{(a-d-f)}\cdot(n-a)_{(b+c-d-e-f)},$$

$$|E_{\alpha_1}^{\widetilde{\alpha_2}\circ\widetilde{\alpha_3}}(n)|=2^{a+b+c-d-2e-2g}\cdot(n-a-b+d)_{(c-e-g)}\cdot(n-c)_{(a+b-d-e-g)}.$$

The products $(\alpha_1*\alpha_2)*\alpha_3$ and $\alpha_1*(\alpha_2*\alpha_3)$ given in equations (10) and (11) as sums over X_1 and X_2 can be written as sums over the set X as follows:

$$\begin{aligned}(\alpha_1*\alpha_2)*\alpha_3=\sum_{(\epsilon_1,\epsilon_2,\epsilon_3)\in X}\quad&\frac{1}{2^{2a+2b+2c-3d-3e-3f}}\cdot\big((n-b-c+e)_{(a-d-f)}\big)^2\\&\cdot(n-a)_{(b+c-d-e-f)}\cdot(n-c)_{(b-e)}\cdot(n-b)_{(c-e)}\\&\epsilon_1\circ\epsilon_2\circ\epsilon_3,\end{aligned}$$

and

$$\begin{aligned}\alpha_1 * (\alpha_2 * \alpha_3) = \sum_{(\epsilon_1,\epsilon_2,\epsilon_3)\in X} \quad & \frac{1}{2^{2a+2b+2c-3d-3e-3g}} \cdot ((n-a-b+d)_{(c-e-g)})^2 \\ & \cdot (n-c)_{(a+b-d-e-g)} \cdot (n-a)_{(b-d)} \cdot (n-b)_{(a-d)} \\ & \epsilon_1 \circ \epsilon_2 \circ \epsilon_3.\end{aligned} \tag{12}$$

For any positive integer n, we have the following easy identities:

$$\begin{array}{rcl}(n-b-c+e)_{(a-d-f)}.(n-c)_{(b-e)} & = & (n-c)_{(a+b-d-e-f)} \\ (n-a)_{(b-d)}.(n-a-b+d)_{(c-e-f)} & = & (n-a)_{(b+c-d-e-f)} \\ (n-b)_{(c-e)}.(n-b-c+e)_{(a-d-f)} & = & (n-b)_{(a-d)}.(n-a-b+d)_{(c-e-f)}.\end{array}$$

Thus, the product $(\alpha_1 * \alpha_2) * \alpha_3$ can be written as follows:

$$\begin{aligned}(\alpha_1 * \alpha_2) * \alpha_3 = \sum_{(\epsilon_1,\epsilon_2,\epsilon_3)\in X} \quad & \frac{1}{2^{2a+2b+2c-3d-3e-3f}} \cdot ((n-a-b+d)_{(c-e-f)})^2 \\ & \cdot (n-c)_{(a+b-d-e-f)} \cdot (n-a)_{(b-d)} \cdot (n-b)_{(a-d)} \\ & \epsilon_1 \circ \epsilon_2 \circ \epsilon_3.\end{aligned} \tag{13}$$

In the set X, the equality $|\delta_2| = |\delta_1|$ can be written $2c+2b-2e+2a-2d-2f = 2a+2b-2d+2c-2e-2g$, so we have $f = g$. Comparison (12) and (13), we see that products $(\alpha_1 * \alpha_2) * \alpha_3$ and $\alpha_1 * (\alpha_2 * \alpha_3)$ are equal, therefore we get the associativity.

3.3. **Action of $\mathcal{B}_n \times \mathcal{B}_n$ on $\mathcal{D}_n$.** In this section, we build the algebra $\mathcal{A}_n$ as the algebra of invariant elements by an action of $\mathcal{B}_n \times \mathcal{B}_n$ on $\mathcal{D}_n$.

Definition 3.3. The group $\mathcal{B}_n \times \mathcal{B}_n$ acts on Q_n by:

$$(a,b) \bullet (\sigma, d, d') = (a\sigma b^{-1}, b(d), a(d')),$$

for any $(a,b) \in \mathcal{B}_n \times \mathcal{B}_n$ and $(\sigma, d, d') \in Q_n$.

Observation. Two partial bijections are in the same orbit if and only if they have the same coset-type.

We can extend this action by linearity to get an action of $\mathcal{B}_n \times \mathcal{B}_n$ on $\mathcal{D}_n$.

Lemma 3.6. *For any three permutations a, b and c of $\mathcal{B}_n$ and for any partial bijections α_1, α_2 of n, the set $E_{\alpha_1}^{\alpha_2}(n)$ is in bijection with $E_{(a,b)\bullet\alpha_1}^{(b,c)\bullet\alpha_2}(n)$.*

Proof. We can check easily that the two following functions:

$$\begin{array}{rccc}\Theta & : & E_{\alpha_1}^{\alpha_2}(n) & \rightarrow & E_{(a,b)\bullet\alpha_1}^{(b,c)\bullet\alpha_2}(n) \\ & & (\widetilde{\alpha_1}, \widetilde{\alpha_2}) & \mapsto & ((a,b)\bullet\widetilde{\alpha_1}, (b,c)\bullet\widetilde{\alpha_2})\end{array},$$

and

$$\begin{array}{rccc}\Psi & : & E_{(a,b)\bullet\alpha_1}^{(b,c)\bullet\alpha_2}(n) & \rightarrow & E_{\alpha_1}^{\alpha_2}(n) \\ & & (\beta_1, \beta_2) & \mapsto & ((a^{-1},b^{-1})\bullet\beta_1, (b^{-1},c^{-1})\bullet\beta_2)\end{array},$$

are well defined. Besides, they are inverse from each other:

$$\begin{array}{rcl}\Psi\Big(\Theta\big((\widetilde{\alpha_1}, \widetilde{\alpha_2})\big)\Big) & = & \Psi\Big(\big((a,b)\bullet\widetilde{\alpha_1}, (b,c)\bullet\widetilde{\alpha_2}\big)\Big) \\ & = & \Big((a^{-1},b^{-1})\bullet(a,b)\bullet\widetilde{\alpha_1}, (b^{-1},c^{-1})\bullet(b,c)\bullet\widetilde{\alpha_2}\Big) \\ & = & (\widetilde{\alpha_1}, \widetilde{\alpha_2}),\end{array}$$

and, similarly,

$$\Theta\Big(\Psi\big(\beta_1,\beta_2\big)\Big) \quad = \quad (\beta_1,\beta_2).$$

Thus Θ defines a bijection between $E_{\alpha_1}^{\alpha_2}(n)$ and $E_{(a,b)\bullet\alpha_1}^{(b,c)\bullet\alpha_2}(n)$ with inverse Ψ. □

It follows from this lemma that the action $\bullet$ is compatible with the product of Q_n. Namely, we can prove the following corollary.

Corollary 3.7. *For any $(a,b,c)\in\mathcal{B}_n^3$ and for any partial bijections α_1,α_2 of n, we have:*

$$(a,c)\bullet(\alpha_1*\alpha_2) = ((a,b)\bullet\alpha_1)*((b,c)\bullet\alpha_2). \tag{14}$$

Proof. If $(\widetilde{\alpha_1},\widetilde{\alpha_2})\in E_{\alpha_1}^{\alpha_2}(n)$, we have:

$$\begin{aligned}
\big((a,b)\bullet\widetilde{\alpha_1}\big)\circ\big((b,c)\bullet\widetilde{\alpha_2}\big) \quad &= \quad \big(a\widetilde{\sigma_1}b^{-1}, b(\widetilde{d_1}), a(\widetilde{d'_1})\big)\circ\big(b\widetilde{\sigma_2}c^{-1}, c(\widetilde{d_2}), b(\widetilde{d'_2})\big)\\
&= \quad (a\widetilde{\sigma_1}\widetilde{\sigma_2}c^{-1}, c(\widetilde{d_2}), a(\widetilde{d'_1}))\\
&= \quad (a,c)\bullet(\widetilde{\alpha_1}\circ\widetilde{\alpha_2}).
\end{aligned}$$

Then, we can write:

$$\begin{aligned}
(a,c)\bullet(\alpha_1*\alpha_2) \quad &= \quad \frac{1}{|E_{\alpha_1}^{\alpha_2}(n)|}\sum_{(\widetilde{\alpha_1},\widetilde{\alpha_2})\in E_{\alpha_1}^{\alpha_2}(n)}(a,c)\bullet(\widetilde{\alpha_1}\circ\widetilde{\alpha_2})\\
&= \quad \frac{1}{|E_{\alpha_1}^{\alpha_2}(n)|}\sum_{(\widetilde{\alpha_1},\widetilde{\alpha_2})\in E_{\alpha_1}^{\alpha_2}(n)}\big((a,b)\bullet\widetilde{\alpha_1}\big)\circ\big((b,c)\bullet\widetilde{\alpha_2}\big)\\
&= \quad \frac{1}{|E_{(a,b)\bullet\alpha_1}^{(b,c)\bullet\alpha_2}(n)|}\sum_{(\widetilde{(a,b)\bullet\alpha_1},\widetilde{(b,c)\bullet\alpha_2})\in E_{(a,b)\bullet\alpha_1}^{(b,c)\bullet\alpha_2}(n)}\widetilde{(a,b)\bullet\alpha_1}\circ\widetilde{(b,c)\bullet\alpha_2}\\
&= \quad ((a,b)\bullet\alpha_1)*((b,c)\bullet\alpha_2)).
\end{aligned}$$

□

We consider the set $\mathcal{A}_n$ of invariant elements by the action of $\mathcal{B}_n\times\mathcal{B}_n$ on $\mathcal{D}_n$:

$$\mathcal{A}_n = \mathcal{D}_n^{\mathcal{B}_n\times\mathcal{B}_n} = \{x\in\mathcal{D}_n \mid (a,b)\bullet x = x \text{ for any } (a,b)\in\mathcal{B}_n\times\mathcal{B}_n\}.$$

Proposition 3.8. *The set $\mathcal{A}_n$ is an algebra with basis the elements $(S_{\lambda,n})_{|\lambda|\leq n}$, where $S_{\lambda,n} = \displaystyle\sum_{\alpha\in Q_n, ct(\alpha)=\lambda}\alpha$.*

Proof. For every $(a,b)\in\mathcal{B}_n\times\mathcal{B}_n$, and for every $x,y\in\mathcal{A}_n$, we have by linearity:

$$(a,b)\bullet(x*y) = ((a,id)\bullet x)*((id,b)\bullet y) = x*y.$$

So $\mathcal{A}_n$ is an algebra.

Any element $x\in\mathcal{D}_n$ writes $x = \displaystyle\sum_{k=1}^{n}\sum_{\substack{d,d'\in\mathbf{P}_n\\|d|=|d'|=2k}}\sum_{\substack{\sigma:d\to d'\\\text{bijection}}} c_{(\sigma,d,d')}(\sigma,d,d')$. If, furthermore x is in $\mathcal{A}_n$, then for every $(a,b)\in\mathcal{B}_n\times\mathcal{B}_n$ we have:

$$\sum_{k=1}^{n}\sum_{\substack{d,d'\in\mathbf{P}_n\\|d|=|d'|=2k}}\sum_{\substack{\sigma:d\to d'\\\text{bijection}}} c_{(\sigma,d,d')}(a\sigma b^{-1}, b(d), a(d')) = \sum_{k=1}^{n}\sum_{\substack{d,d'\in\mathbf{P}_n\\|d|=|d'|=2k}}\sum_{\substack{\sigma:d\to d'\\\text{bijection}}} c_{(\sigma,d,d')}(\sigma,d,d').$$

Thus, for any $(a,b) \in \mathcal{B}_n \times \mathcal{B}_n$, we have $c_{(a\sigma b^{-1}, b(d), a(d'))} = c_{(\sigma, d, d')}$. This means that if $x \in \mathcal{A}_n$, all partial permutations in the same orbit – that is with the same coset-type – have the same coefficients. Therefore, the elements $(S_{\lambda,n})_{|\lambda| \leq n}$ form a basis of $\mathcal{A}_n$. □

Corollary 3.9. *If λ and δ are two partitions such that $|\lambda|, |\delta| \leq n$, there exist unique constants $c_{\lambda\delta}^{\rho}(n) \in \mathbb{C}$ such that:*

$$\mathcal{S}_{\lambda,n} * \mathcal{S}_{\delta,n} = \sum_{\substack{\rho \text{ partition} \\ \max(|\lambda|,|\delta|) \leq |\rho| \leq \min(|\lambda|+|\delta|, n)}} c_{\lambda\delta}^{\rho}(n) \mathcal{S}_{\rho,n}.$$

Proof. We only have to prove the inequalities on the size of ρ. Let α_1 and α_2 be two partial bijections of n with coset-type λ and δ. By definition (see Figure 2), every partial bijection of n that appears in the sum of the product $\alpha_1 * \alpha_2$ has some coset-type ρ with $|\rho| = \frac{|d_1 \cup d_2'|}{2}$. But

$$\max\left(\frac{|d_1|}{2}, \frac{|d_2'|}{2}\right) = \max(|\lambda|, |\delta|) \leq |\rho| = \frac{|d_1 \cup d_2'|}{2} \leq \frac{|d_1| + |d_2'|}{2} = |\lambda| + |\delta|. \quad \square$$

Lemma 3.10. *Let λ be a partition such that $|\lambda| = r \leq n$, we have:*

$$\psi_n(\mathcal{S}_{\lambda,n}) = \frac{1}{2^{n-|\lambda|}(n-|\lambda|)!} \binom{n-|\bar{\lambda}|}{m_1(\lambda)} K_{\bar{\lambda}}(n).$$

Proof. We first prove the following equation:

$$\sum_{\alpha \in A_{\lambda,n}} \sum_{\hat{\alpha} \in \mathcal{S}_{2n} \cap P_\alpha(n)} \hat{\sigma} = \binom{n-|\bar{\lambda}|}{m_1(\lambda)} K_{\bar{\lambda}}(n). \tag{15}$$

Fix a permutation $\omega \in K_{\bar{\lambda}}(n)$, that is $\omega \in \mathcal{S}_{2n}$ and $ct(\omega) = \bar{\lambda} \cup 1^{n-|\bar{\lambda}|}$. We are looking for the number of partial bijections $\alpha \in A_{\lambda,n}$ such that ω is one of its trivial extensions. We denote by $\mathrm{supp}(\omega)$ the set $\mathrm{supp}(\omega) \subseteq d$ such that $ct(\omega_{|_{\mathrm{supp}(\omega)}}) = \bar{\lambda}$. The following condition is necessary so that ω is a trivial extension of α: $\alpha_{|_{\mathrm{supp}(\omega)}}$ must be equal to $\omega_{|_{\mathrm{supp}(\omega)}}$. Thus the partial bijections α we are looking for are the restrictions of ω to sets $\mathrm{supp}(\omega) \cup x$, where $|\mathrm{supp}(\omega) \cup x| = 2|\lambda|$. Since $|\mathrm{supp}(\omega)| = 2|\bar{\lambda}|$, their number is $\binom{n-|\bar{\lambda}|}{|\lambda|-|\bar{\lambda}|} = \binom{n-|\bar{\lambda}|}{m_1(\lambda)}$. This ends the proof of equation (15).
By applying ψ_n to $\mathcal{S}_{\lambda,n}$, we get:

$$\begin{aligned} \psi_n(\mathcal{S}_{\lambda,n}) &= \psi_n\Big(\sum_{\alpha \in A_{\lambda,n}} \alpha\Big) \\ &= \frac{1}{2^{n-|\lambda|}(n-|\lambda|)!} \sum_{\alpha \in A_{\lambda,n}} \sum_{\hat{\alpha} \in \mathcal{S}_{2n} \cap P_\alpha(n)} \hat{\sigma} \\ &= \frac{1}{2^{n-|\lambda|}(n-|\lambda|)!} \binom{n-|\bar{\lambda}|}{m_1(\lambda)} K_{\bar{\lambda}}(n). \end{aligned}$$

□

This lemma implies that $\psi_n(\mathcal{A}_n) \subseteq \mathbb{C}[\mathcal{B}_n/\mathcal{S}_{2n}\backslash\mathcal{B}_n]$. The morphism $\mathcal{A}_n \rightarrow \mathbb{C}[\mathcal{B}_n/\mathcal{S}_{2n}\backslash\mathcal{B}_n]$ mentioned in Section 2.5 is the morphism $\psi_{n_{|_{\mathcal{A}_n}}}$.

3.4. **Morphism from $\mathcal{A}_{n+1}$ to $\mathcal{A}_n$.** This paragraph is dedicated to prove the following proposition:

Proposition 3.11. *The function φ_n defined as follows:*

$$\begin{array}{cccc} \varphi_n & : & \mathcal{A}_{n+1} & \rightarrow & \mathcal{A}_n \\ & & \mathcal{S}_{\lambda,n+1} & \mapsto & \begin{cases} \frac{n+1}{(n+1-|\lambda|)} \mathcal{S}_{\lambda,n} & \text{if } \quad |\lambda| < n+1, \\ 0 & \text{if } \quad |\lambda| = n+1, \end{cases} \end{array}$$

is a morphism of algebras.

Let $\mathcal{S}_{\lambda,n+1}$ and $\mathcal{S}_{\delta,n+1}$ where $|\lambda| \leq n+1$ and $|\delta| \leq n+1$ be two basis elements of $\mathcal{A}_{n+1}$.

If λ (resp. δ) is a partition of $n+1$, then $\varphi_n(\mathcal{S}_{\lambda,n+1})$ (resp. $\varphi_n(\mathcal{S}_{\delta,n+1})$) is equal to zero, and by Corollary 3.9 we have:

$$\mathcal{S}_{\lambda,n+1} * \mathcal{S}_{\delta,n+1} = \sum_{\substack{\rho \text{ partition} \\ |\rho|=n+1}} c_{\lambda\delta}^{\rho}(n+1)\mathcal{S}_{\rho,n+1}.$$

The size of partitions ρ indexing the sum in this equation is $n+1$. By applying φ_n, we get:

$$\varphi_n(\mathcal{S}_{\lambda,n+1} * \mathcal{S}_{\delta,n+1}) = \sum_{\substack{\rho \text{ partition} \\ |\rho|=n+1}} c_{\lambda\delta}^{\rho}(n+1)\varphi_n(\mathcal{S}_{\rho,n+1}) = 0.$$

Thus in this case we have $\varphi_n(\mathcal{S}_{\lambda,n+1} * \mathcal{S}_{\delta,n+1}) = \varphi_n(\mathcal{S}_{\lambda,n+1}) * \varphi_n(\mathcal{S}_{\delta,n+1})$.

In the other case ($|\lambda| \leq n$ and $|\delta| \leq n$) we have by Corollary 3.9:

$$\mathcal{S}_{\lambda,n+1} * \mathcal{S}_{\delta,n+1} = \sum_{\substack{r \leq n+1 \\ \rho \vdash r}} c_{\lambda\delta}^{\rho}(n+1)\mathcal{S}_{\rho,n+1}.$$

This gives us the following equation after applying φ_n:

$$\begin{aligned} \varphi_n(\mathcal{S}_{\lambda,n+1} * \mathcal{S}_{\delta,n+1}) &= \varphi_n\left(\sum_{\substack{r \leq n+1 \\ \rho \vdash r}} c_{\lambda\delta}^{\rho}(n+1)\mathcal{S}_{\rho,n+1}\right) \\ &= \sum_{\substack{r \leq n \\ \rho \vdash r}} c_{\lambda\delta}^{\rho}(n+1)\frac{n+1}{(n+1-|\rho|)}\mathcal{S}_{\rho,n}. \end{aligned}$$

In the other hand, we have:

$$\begin{aligned} \varphi_n(\mathcal{S}_{\lambda,n+1}) * \varphi_n(\mathcal{S}_{\delta,n+1}) &= \frac{n+1}{(n+1-|\lambda|)}\mathcal{S}_{\lambda,n} * \frac{n+1}{(n+1-|\delta|)}\mathcal{S}_{\delta,n} \\ &= \frac{n+1}{(n+1-|\lambda|)}\frac{n+1}{(n+1-|\delta|)}\sum_{\substack{r \leq n \\ \rho \vdash r}} c_{\lambda\delta}^{\rho}(n)\mathcal{S}_{\rho,n}. \end{aligned}$$

Thus, φ_n is a morphism if we have the following equality for any partition ρ with size less than n:

$$\frac{c_{\lambda\delta}^{\rho}(n+1)}{c_{\lambda\delta}^{\rho}(n)} = \frac{\frac{n+1}{(n+1-|\lambda|)}\frac{n+1}{(n+1-|\delta|)}}{\frac{n+1}{(n+1-|\rho|)}}.$$

Let ρ be a partition with size less then n and α an element of $A_{\rho,n}$. We define $H_{\lambda\delta}^{\rho}(n)$ to be the following set:

$$\{(\alpha_1, \alpha_2) \in A_{\lambda,n} \times A_{\delta,n} \text{ such that there exists } (\widetilde{\alpha_1}, \widetilde{\alpha_2}) \in E_{\alpha_1}^{\alpha_2}(n) \text{ with } \alpha = \widetilde{\alpha_1} \circ \widetilde{\alpha_2}\}.$$

This set depends on α by definition. However, α does not appear in our notation. This should not be an issue, since α is fixed in the whole proof.

The coefficient $c_{\lambda\delta}^{\rho}(n)$ can be written as follows:

$$c_{\lambda\delta}^{\rho}(n) = \sum_{(\alpha_1,\alpha_2) \in H_{\lambda\delta}^{\rho}(n)} \frac{1}{|E_{\alpha_1}^{\alpha_2}(n)|}.$$

Similarly, we have:

$$c^{\rho}_{\lambda\delta}(n+1) = \sum_{(\alpha_1,\alpha_2)\in H^{\rho}_{\lambda\delta}(n+1)} \frac{1}{|E^{\alpha_2}_{\alpha_1}(n+1)|}.$$

By equation (4), if $(\alpha_1, \alpha_2) \in H^{\rho}_{\lambda,\delta}(n)$, we have:

$$|E^{\alpha_2}_{\alpha_1}(n)| = 2^{2|\rho|-|\lambda|-|\delta|}(n-|\lambda|)_{(|\rho|-|\lambda|)}(n-|\delta|)_{(|\rho|-|\delta|)}.$$

Similarly, if $(\alpha_1, \alpha_2) \in H^{\rho}_{\lambda,\delta}(n+1)$, we have:

$$|E^{\alpha_2}_{\alpha_1}(n+1)| = 2^{2|\rho|-|\lambda|-|\delta|}(n+1-|\lambda|)_{(|\rho|-|\lambda|)}(n+1-|\delta|)_{(|\rho|-|\delta|)}.$$

Thus, we get:

$$c^{\rho}_{\lambda\delta}(n) = \frac{|H^{\rho}_{\lambda\delta}(n)|}{2^{2|\rho|-|\lambda|-|\delta|}(n-|\lambda|)_{(|\rho|-|\lambda|)}(n-|\delta|)_{(|\rho|-|\delta|)}},$$

and

$$c^{\rho}_{\lambda\delta}(n+1) = \frac{|H^{\rho}_{\lambda\delta}(n+1)|}{2^{2|\rho|-|\lambda|-|\delta|}(n+1-|\lambda|)_{(|\rho|-|\lambda|)}(n+1-|\delta|)_{(|\rho|-|\delta|)}}.$$

This gives us after simplification:

$$\frac{c^{\rho}_{\lambda\delta}(n+1)}{c^{\rho}_{\lambda\delta}(n)} = \frac{|H^{\rho}_{\lambda\delta}(n+1)|}{|H^{\rho}_{\lambda\delta}(n)|} \cdot \frac{n+1-|\rho|}{n+1-|\lambda|} \cdot \frac{n+1-|\rho|}{n+1-|\delta|}$$

We will now evaluate the quotient $\frac{|H^{\rho}_{\lambda\delta}(n+1)|}{|H^{\rho}_{\lambda\delta}(n)|}$. Let $\boldsymbol{u} = (u_1, u'_1, u_2, u'_2)$ be an element of $\mathbf{P}^4_n$ such that:

(16) $$u_2 \subseteq d,$$

(17) $$u'_1 \subseteq d',$$

(18) $$|u_1| = |u'_1| = 2|\lambda|,$$

(19) $$|u_2| = |u'_2| = 2|\delta|,$$

(20) $$|u'_2 \cup u_1| = 2|\rho|.$$

We denote by

$$\begin{aligned} N_{\boldsymbol{u}} \quad = \quad & \{\big(h_1 = (f_1, u_1, u'_1), h_2 = (f_2, u_2, u'_2)\big) \in A_{\lambda,n} \times A_{\delta,n} \\ & \text{such that there exists } (\widetilde{h_1}, \widetilde{h_2}) \in E^{h_2}_{h_1}(n) \text{ with } \alpha = \widetilde{h_1} \circ \widetilde{h_2}\}. \end{aligned}$$

Its elements are represented on Figure 6. The set $H^{\rho}_{\lambda\delta}(n)$ is the disjoint union of all $N_{\boldsymbol{u}}$ with $\boldsymbol{u}$ satisfying the above conditions.

Lemma 3.12. *Let $\boldsymbol{v} = (v_1, v'_1, v_2, v'_2)$ be an element of $\mathbf{P}^4_n$ satisfying conditions above. If $v'_1 = u'_1$ and $v_2 = u_2$, then there exists a bijection between $N_{\boldsymbol{u}}$ and $N_{\boldsymbol{v}}$.*

Proof. We take any permutation $b \in \mathcal{B}_n$, such that $b(u_1) = v_1$ and $b(u'_2) = v'_2$. Such a permutation exists because $|u'_2 \cup u_1| = |v'_2 \cup v_1|$. We associate to a pair (h_1, h_2) in $N_{\boldsymbol{u}}$ the pair $((id, b) \bullet h_1, (b, id) \bullet h_2)$. We check that the image lies in $N_{\boldsymbol{v}}$:

$$\begin{aligned} \big((id, b) \bullet h_1, (b, id) \bullet h_2\big) &= \big((f_1 b^{-1}, b(u_1), u'_1), (b f_2, u_2, b(u'_2))\big) \\ &= \big((f_1 b^{-1}, v_1, u'_1), (b f_2, u_2, v'_2)\big) \in A_{\lambda,n} \times A_{\delta,n}. \end{aligned}$$



FIGURE 6. Schematic representation of elements of $N_{\boldsymbol{u}}$

We can check easily that $\big((id,b)\bullet\widetilde{h_1},(b,id)\bullet\widetilde{h_2}\big)\in E^{(b,id)\bullet h_2}_{(id,b)\bullet h_1}(n)$, and we have:

$$\begin{aligned}(id,b)\bullet\widetilde{h_1}\circ(b,id)\bullet\widetilde{h_2} &= (\widetilde{f_1}b^{-1},b(\widetilde{u_1}),\widetilde{u'_1})\circ(b\widetilde{f_2},\widetilde{u_2},b(\widetilde{u'_2}))\\ &= (\widetilde{f_1}b^{-1}b\widetilde{f_2},\widetilde{u_2},\widetilde{u'_1})\\ &= \widetilde{h_1}\circ\widetilde{h_2}\\ &= \alpha.\end{aligned}$$

It is then easy to check that this defines a bijection between $N_{\boldsymbol{u}}$ and $N_{\boldsymbol{v}}$. Details are similar to Lemma 3.6's proof. □

Therefore the cardinality of $N_{\boldsymbol{u}}$ depends only on u'_1 and u_2. We denote it by $f(u'_1,u_2)$. If we denote by $\boldsymbol{U}$ the set of vectors $\boldsymbol{u}\in\mathbf{P}^4_n$ satisfying conditions (16) to (20), the set $H^\rho_{\lambda\delta}(n)$ can be written as follows:

$$H^\rho_{\lambda\delta}(n)=\bigsqcup_{\boldsymbol{u}\in\boldsymbol{U}}N_{\boldsymbol{u}}.$$

Using Lemma 3.12, we obtain:

$$|H^\rho_{\lambda\delta}(n)|=\sum_{\boldsymbol{u}\in\boldsymbol{U}}|N_{\boldsymbol{u}}|=\sum_{u'_1,u_2}\sum_{u_1,u'_2}f(u'_1,u_2).$$

The first (resp. second) summation indexes are vectors u'_1 and u_2 (resp. u_1 and u'_2) satisfying conditions (16) to (19) (resp. (18) to (20)). Since $N_{\boldsymbol{u}}$ depends only on u'_1 and u_2, we get:

$$|H^\rho_{\lambda\delta}(n)|=\sum_{u'_1,u_2}f(u'_1,u_2)k_n,$$

where k_n is the number of possible choices of vectors u_1 and u'_2 satisfying conditions (18) to (20). There are $\binom{n}{|\lambda|}$ sets $u_1\in\mathbf{P}_n$ that fulfill (18). Once u_1 is chosen, it remains $\binom{|\lambda|}{|\lambda|+|\delta|-|\rho|}\cdot\binom{n-|\lambda|}{|\rho|-|\lambda|}$ ways to choose u'_2 with conditions (19) and (20). The first binomial is the number of possible choices of $u_1\cap u'_2$ and the second one is the number of possible choices of $u'_2\setminus u_1$. Then, we have:

$$k_n=\binom{n}{|\lambda|}\cdot\binom{n-|\lambda|}{|\rho|-|\lambda|}\cdot\binom{|\lambda|}{|\lambda|+|\delta|-|\rho|}.$$

Thus, the cardinality of $H^{\rho}_{\lambda\delta}(n)$ is:

$$|H^{\rho}_{\lambda\delta}(n)| = \binom{n}{|\lambda|} \cdot \binom{n-|\lambda|}{|\rho|-|\lambda|} \cdot \binom{|\lambda|}{|\lambda|+|\delta|-|\rho|} \sum_{u_1', u_2} f(u_1', u_2).$$

The summation index does not depend on n because u_1' and u_2 should fulfill conditions (16) and (17). Similarly, we obtain:

$$|H^{\rho}_{\lambda\delta}(n+1)| = \binom{n+1}{|\lambda|} \cdot \binom{n+1-|\lambda|}{|\rho|-|\lambda|} \cdot \binom{|\lambda|}{|\lambda|+|\delta|-|\rho|} \sum_{u_1', u_2} f(u_1', u_2),$$

which gives us:

$$\frac{|H^{\rho}_{\lambda\delta}(n+1)|}{|H^{\rho}_{\lambda\delta}(n)|} = \frac{\binom{n+1}{|\delta|}\binom{n+1-|\delta|}{|\rho|-|\delta|}}{\binom{n}{|\delta|}\binom{n-|\delta|}{|\rho|-|\delta|}} = \frac{n+1}{n+1-|\rho|}.$$

Thus, we have:

$$\begin{aligned} \frac{c^{\rho}_{\lambda\delta}(n+1)}{c^{\rho}_{\lambda\delta}(n)} &= \frac{n+1}{n+1-|\rho|} \cdot \frac{n+1-|\rho|}{n+1-|\lambda|} \cdot \frac{n+1-|\rho|}{n+1-|\delta|} \\ &= \frac{\frac{n+1}{(n+1-|\lambda|)}\frac{n+1}{(n+1-|\delta|)}}{\frac{n+1}{(n+1-|\rho|)}}. \end{aligned} \tag{21}$$

This proves that φ_n is a morphism of algebras.

3.5. **Projective limits.** In this paragraph, we consider the projective limit $\mathcal{A}_\infty$ of the sequence $(\mathcal{A}_n)$. We prove in Proposition 3.15 that every element of $\mathcal{A}_\infty$ is written in a unique way as infinite linear combination of elements indexed by partitions.

First, from equation (21), we can get the following Corollary:

Corollary 3.13. *Let λ, δ and ρ be three partitions such that*

$$\max(|\lambda|, |\delta|) \leq |\rho| \leq |\lambda| + |\delta|.$$

For every $n \geq |\rho|$, we have:

$$c^{\rho}_{\lambda\delta}(n) = \frac{c^{\rho}_{\lambda\delta}(|\rho|)}{\binom{|\rho|}{|\lambda|}\binom{|\rho|}{|\delta|}} \cdot \frac{\binom{n}{|\lambda|}\binom{n}{|\delta|}}{\binom{n}{|\rho|}}.$$

Proof. We proceed by induction on n. For $n = |\rho|$, we have the equality. Assume we have the equality for some $n \geq |\rho|$ and let us prove it for $n+1$. By equation (21), we have:

$$\frac{c^{\rho}_{\lambda\delta}(n+1)}{c^{\rho}_{\lambda\delta}(n)} = \frac{\frac{n+1}{(n+1-|\lambda|)}\frac{n+1}{(n+1-|\delta|)}}{\frac{n+1}{(n+1-|\rho|)}}.$$

This gives us the following equality, using the induction hypothesis :

$$\begin{aligned} c^{\rho}_{\lambda\delta}(n+1) &= \frac{c^{\rho}_{\lambda\delta}(|\rho|)}{\binom{|\rho|}{|\lambda|}\binom{|\rho|}{|\delta|}} \cdot \frac{\binom{n}{|\lambda|}\binom{n}{|\delta|}}{\binom{n}{|\rho|}} \cdot \frac{\frac{n+1}{(n+1-|\lambda|)}\frac{n+1}{(n+1-|\delta|)}}{\frac{n+1}{(n+1-|\rho|)}} \\ &= \frac{c^{\rho}_{\lambda\delta}(|\rho|)}{\binom{|\rho|}{|\lambda|}\binom{|\rho|}{|\delta|}} \cdot \frac{\binom{n+1}{|\lambda|}\binom{n+1}{|\delta|}}{\binom{n+1}{|\rho|}}. \end{aligned}$$

□

Let $\mathcal{A}_\infty$ be the projective limit of $(\mathcal{A}_n, \varphi_n)$:

$$\mathcal{A}_\infty = \{(a_n)_{n\geq 1} \mid \text{for every } n \geq 1, a_n \in \mathcal{A}_n \text{ and } \varphi_n(a_{n+1}) = a_n\}.$$

Lemma 3.14. *An element $a = (a_n)_{n\geq 1}$ is in $\mathcal{A}_\infty$ if and only if there exists a family $(x^a_\lambda)_{\lambda\ partition}$ of elements of $\mathbb{C}$ such that for every $n \geq 1$, $a_n = \sum_{\substack{\lambda\ partition \\ |\lambda|\leq n}} \frac{x^a_\lambda}{\binom{n}{|\lambda|}} S_{\lambda,n}$.*

Proof. Let $a = (a_n)_{n\geq 1}$ be a series in $\mathcal{A}_\infty$, $a_n \in \mathcal{A}_n$ for every $n \geq 1$. By the Proposition 3.8, the elements $(\mathcal{S}_{\lambda,n})_{\lambda\vdash r\leq n}$ form a basis of $\mathcal{A}_n$, thus for every $n \geq 1$ and every partition λ such as $|\lambda| \leq n$, there exists a scalar $a_\lambda(n) \in \mathbb{C}$ such that

$$a_n = \sum_{\substack{\lambda \text{ partition} \\ |\lambda|\leq n}} a_\lambda(n)\mathcal{S}_{\lambda,n}.$$

The condition $\varphi_n(a_{n+1}) = a_n$ can be written as follows:

$$\varphi_n\left(\sum_{\substack{\lambda \text{ partition} \\ |\lambda|\leq n+1}} a_\lambda(n+1)\mathcal{S}_{\lambda,n+1}\right) = \sum_{\substack{\lambda \text{ partition} \\ |\lambda|\leq n}} a_\lambda(n)\mathcal{S}_{\lambda,n}.$$

Using the definition of φ_n, we can simplify this equality to obtain:

$$\sum_{\substack{\lambda \text{ partition} \\ |\lambda|\leq n}} a_\lambda(n+1)\frac{n+1}{n+1-|\lambda|}\mathcal{S}_{\lambda,n} = \sum_{\substack{\lambda \text{ partition} \\ |\lambda|\leq n}} a_\lambda(n)\mathcal{S}_{\lambda,n}.$$

By considering the coefficients of $\mathcal{S}_{\lambda,n}$ we get that for every partition λ such that $|\lambda| \leq n$, we have:

$$\frac{a_\lambda(n+1)}{a_\lambda(n)} = \frac{n+1-|\lambda|}{n+1}.$$

After an immediate induction, we get :

$$a_\lambda(n) = \frac{a_\lambda(|\lambda|)}{\binom{n}{|\lambda|}}.$$

Set $x^a_\lambda = a_\lambda(|\lambda|)$, this proves the "only if" statement. Converse is obvious. □

For every partition λ, we define the sequence T_λ as follows:

$$T_\lambda = (T_\lambda)_{n\geq 1} = \begin{cases} 0 & \text{if} \quad n < |\lambda|, \\ \frac{1}{\binom{n}{|\lambda|}} S_{\lambda,n} & \text{if} \quad n \geq |\lambda|. \end{cases}$$

From Lemma 3.14, we can directly obtain the following proposition:

Proposition 3.15. *Every element $a \in \mathcal{A}_\infty$ is written in a unique way as infinite linear combination of elements T_λ.*

This proposition shows that the algebra $\mathcal{A}_\infty$ satisfies the second property required in Section 2.5. In particular, $T_\lambda * T_\delta$ writes as linear combination of elements T_ρ. We can be more precise.

Corollary 3.16. *Let λ and δ be two partitions, there exist unique constants $b_{\lambda\delta}^{\rho}$ such that:*

$$T_\lambda * T_\delta = \sum_{\substack{\rho \text{ partition} \\ \max(|\lambda|,|\delta|)\leq|\rho|\leq|\lambda|+|\delta|}} b_{\lambda\delta}^{\rho} T_\rho .$$

Moreover $b_{\lambda\delta}^{\rho} = \frac{c_{\lambda\delta}^{\rho}(|\rho|)}{\binom{|\rho|}{|\lambda|}\binom{|\rho|}{|\delta|}}$. In particular, it is a non-negative rational number.

Proof. By Proposition 3.15, $T_\lambda * T_\delta$ writes as linear combination of elements T_ρ. It remains to prove how we get the conditions about the size of partitions ρ that appear in the sum index and the formula of $b_{\lambda\delta}^{\rho}$.
If $n < \max(|\lambda|, |\delta|)$, we have:

$$(T_\lambda * T_\delta)_n = 0.$$

Let $n \geq \max(|\lambda|, |\delta|)$, we use Corollary 3.9 and Corollary 3.13 to get:

$$\begin{aligned} (T_\lambda * T_\delta)_n &= \frac{1}{\binom{n}{|\lambda|}} S_{\lambda,n} * \frac{1}{\binom{n}{|\delta|}} S_{\delta,n} \\ &= \sum_{\substack{\rho \text{ partition} \\ \max(|\lambda|,|\delta|)\leq|\rho|\leq\min(|\lambda|+|\delta|,n)}} \frac{c_{\lambda\delta}^{\rho}(|\rho|)}{\binom{|\rho|}{|\lambda|}\binom{|\rho|}{|\delta|}\binom{n}{|\rho|}} \mathcal{S}_{\rho,n} \\ &= \left(\sum_{\substack{\rho \text{ partition} \\ \max(|\lambda|,|\delta|)\leq|\rho|\leq|\lambda|+|\delta|}} \frac{c_{\lambda\delta}^{\rho}(|\rho|)}{\binom{|\rho|}{|\lambda|}\binom{|\rho|}{|\delta|}} T_\rho \right)_n \\ &= \left(\sum_{\substack{\rho \text{ partition} \\ \max(|\lambda|,|\delta|)\leq|\rho|\leq|\lambda|+|\delta|}} b_{\lambda\delta}^{\rho} T_\rho \right)_n . \end{aligned}$$

This proves our proposition. ☐

Example 3.1. *We compute in this example the product* $T_{(2)} * T_{(2)}$. *Using Corollary 3.16, we can write,* $T_{(2)} * T_{(2)} = \sum_{\substack{\rho \textit{ partition} \\ 2\leq|\rho|\leq 4}} b^{\rho}_{(2)(2)} T_\rho$, *which gives us:*

$$\begin{aligned} T_{(2)} * T_{(2)} &= b^{(1^2)}_{(2)(2)} T_{(1^2)} + b^{(1^3)}_{(2)(2)} T_{(1^3)} + b^{(1^4)}_{(2)(2)} T_{(1^4)} \\ &\quad + b^{(2)}_{(2)(2)} T_{(2)} + b^{(2,1)}_{(2)(2)} T_{(2,1)} + b^{(2,1^2)}_{(2)(2)} T_{(2,1^2)} + b^{(2^2)}_{(2)(2)} T_{(2^2)} \\ &\quad + b^{(3)}_{(2)(2)} T_{(3)} + b^{(3,1)}_{(2)(2)} T_{(3,1)} \\ &\quad + b^{(4)}_{(2)(2)} T_{(4)}. \end{aligned}$$

The formula for $b^\rho_{\lambda\delta}$ *given in Corollary 3.16 shows that these elements can be computed using the product of* $\mathcal{S}_{(2),|\lambda|+|\delta|} * \mathcal{S}_{(2),|\lambda|+|\delta|}$ *in* $\mathcal{A}_{|\lambda|+|\delta|}$, *which is* $\mathcal{A}_4$ *in our case.*
Using Sage *program* [Sag], *we get the following equation for the product* $\mathcal{S}_{(2),4} * \mathcal{S}_{(2),4}$ *in* $\mathcal{A}_4$*:*

$$\mathcal{S}_{(2),4} * \mathcal{S}_{(2),4} = 96\mathcal{S}_{(1^2),4} + 48\mathcal{S}_{(2),4} + 36\mathcal{S}_{(3),4} + 12\mathcal{S}_{(2^2),4}.$$

Using formulas of $c^\rho_{\lambda\delta}(|\rho|)$ *given in Corollary 3.13 and* $b^\rho_{\lambda\delta}$, *we obtain:*

$$T_{(2)} * T_{(2)} = 16T_{(1^2)} + 8T_{(2)} + 4T_{(3)} + \frac{1}{3}T_{(2^2)}.$$

4. PROOF OF THEOREM 2.1

In the previous section, we built all algebras and morphisms that we need in order to prove Theorem 2.1.

Let λ and δ be two proper partitions, by Corollary 3.16, we have:

$$T_\lambda * T_\delta = \sum_{\substack{\rho \text{ partition} \\ \max(|\lambda|,|\delta|)\leq|\rho|\leq|\lambda|+|\delta|}} b^\rho_{\lambda\delta} T_\rho.$$

Recall that this is an equality of sequences. Taking the n-th term, we have:

$$\frac{1}{\binom{n}{|\lambda|}} S_{\lambda,n} * \frac{1}{\binom{n}{|\delta|}} S_{\delta,n} = \sum_{\substack{\rho \text{ partition} \\ \max(|\lambda|,|\delta|)\leq|\rho|\leq\min(|\lambda|+|\delta|,n)}} b^\rho_{\lambda\delta} \frac{1}{\binom{n}{|\rho|}} S_{\rho,n}.$$

By applying ψ_n we obtain (see Lemma 3.10):

$$\frac{1}{2^{n-|\lambda|}(n-|\lambda|)!} K_\lambda(n) \cdot \frac{1}{2^{n-|\delta|}(n-|\delta|)!} K_\delta(n) =$$

$$\sum_{\substack{\rho \text{ partition} \\ \max(|\lambda|,|\delta|)\leq|\rho|\leq\min(|\lambda|+|\delta|,n)}} b^\rho_{\lambda\delta} \frac{\binom{n}{|\lambda|}\binom{n}{|\delta|}}{\binom{n}{|\rho|} 2^{n-|\rho|}(n-|\rho|)!} \binom{n-|\bar{\rho}|}{m_1(\rho)} K_{\bar{\rho}}(n).$$

After simplification, we get:

$$K_\lambda(n) \cdot K_\delta(n) = \sum_{\substack{\rho \text{ partition} \\ \max(|\lambda|,|\delta|)\leq|\rho|\leq\min(|\lambda|+|\delta|,n)}} b^\rho_{\lambda\delta} \frac{(|\rho|)_{|\bar{\rho}|}}{|\lambda|!|\delta|!} 2^{n+|\rho|-|\lambda|-|\delta|} n!(n - |\bar{\rho}|)_{m_1(\rho)} K_{\bar{\rho}}(n).$$

Fact. Any partition ρ such that $|\rho| \leq \min(|\lambda|+|\delta|, n)$ can be written in a unique way as $\rho = \tau \cup (1^j)$, where τ is a proper partition and $j \leq \min(|\lambda|+|\delta|, n) - |\tau|$.

Using this fact, the product can be written as follows:

$$K_\lambda(n) \cdot K_\delta(n) = \sum_{\substack{\tau \text{ proper partition} \\ |\tau| \leq \min(|\lambda|+|\delta|, n)}} \alpha_{\lambda\delta}^{\tau}(n) K_\tau(n),$$

where

$$\begin{aligned} \alpha_{\lambda\delta}^{\tau}(n) &= \frac{1}{|\lambda|!|\delta|!} \sum_{j=0}^{\min(|\lambda|+|\delta|,n)-|\tau|} b_{\lambda\delta}^{\tau\cup(1^j)} n!(n-|\tau|)_j(|\tau|+j)_{|\tau|} 2^{n+|\tau|+j-|\lambda|-|\delta|} \\ &= \frac{2^n n!}{|\lambda|!|\delta|!} \sum_{j=0}^{|\lambda|+|\delta|-|\tau|} b_{\lambda\delta}^{\tau\cup(1^j)} (n-|\tau|)_j(|\tau|+j)_{|\tau|} 2^{|\tau|+j-|\lambda|-|\delta|}. \end{aligned} \tag{22}$$

The change of sum index in the last equality comes from the fact that if $n < |\lambda| + |\delta|$, we have:

$$(n-|\tau|)_j = 0 \text{ for any } j \text{ with } n-|\tau| < j \leq |\lambda|+|\delta|-|\tau|.$$

This ends the proof of Theorem 2.1.

Corollary 4.1. *If λ, δ and ρ are three proper partitions such that $|\rho| = |\lambda| + |\delta|$, then:*

$$\alpha_{\lambda\delta}^{\rho}(n) = b_{\lambda\delta}^{\rho} \frac{|\rho|!}{|\lambda|!|\delta|!} 2^n n! = c_{\lambda\delta}^{\rho}(|\rho|) \frac{|\lambda|!|\delta|!}{(|\lambda|+|\delta|)!} 2^n n!.$$

Example 4.1. *We recall that the product $T_{(2)} * T_{(2)}$ has been computed in Example 3.1. We deduce from it the complete formula for the product $K_{(2)}(n) \cdot K_{(2)}(n)$ for every $n \geq 4$. Using formula* (22)*, we have:*

$$K_{(2)}(n) \cdot K_{(2)}(n) = 2^n n! n(n-1) K_\emptyset(n) + 2^n n! K_{(2)}(n) + 2^n n! 3 K_{(3)}(n) + 2^n n! 2 K_{(2^2)}(n).$$

REFERENCES

[AC10] Kürşat Aker and Mahir Bilen Can. Hecke ring of (S_{2n}, H_n): Generators and the Farahat-Higman ring. arXiv preprint 1009.5373, 2010.

[Cor75] R. Cori. Un code pour les graphes planaires et ses applications. Number 27 in Astérisque. Société Mathématique de France, 1975. 169 pages.

[DF12] M. Dołęga and V. Féray. On Kerov polynomials for Jack characters. preprint arXiv:1201.1806, 2012.

[FH59] H. Farahat and G. Higman. The centres of symmetric group rings. *Proc. Roy. Soc. (A)*, 250:212–221, 1959.

[GJ96] I. P. Goulden and D. M. Jackson. Maps in locally orientable surfaces, the double coset algebra, and zonal polynomials. *Can. J. Math.*, 48(3):569–584, 1996.

[GS98] A. Goupil and G. Schaeffer. Factoring n-cycles and counting maps of given genus. *Eur. J. Comb.*, 19(7):819–834, 1998.

[IK99] V. Ivanov and S. Kerov. The algebra of conjugacy classes in symmetric groups, and partial permutations. *Zap. Nauchn. Sem. S.-Peterburg. Otdel. Mat. Inst. Steklov. (POMI)*, 256(3):95–120, 1999.

[Jam61] Alan T. James. Zonal polynomials of the real positive definite symmetric matrices. *Annals of Mathematics*, 74(3):456–469, 1961.

[JV90] D.M. Jackson and T.I. Visentin. A character theoretic approach to embeddings of rooted maps in an orientable surface of given genus. *Trans. AMS*, 322:343–363, 1990.

[Mac95] I.G. Macdonald. *Symmetric functions and Hall polynomials*. Oxford Univ. Press, second edition, 1995.

[Sag] Sage mathematical software, version 2.6, http://www.sagemath.org.

LaBRI, UNIVERSITÉ BORDEAUX 1, 351 COURS DE LA LIBÉRATION, 33 400 TALENCE, FRANCE
E-mail address: `omar.tout@labri.fr`